

Annex 2.

Dear parent!

I would like your consent for the next (re)vaccination of your child (first name and surname) _____ against **diphtheria and tetanus** in accordance with the national immunisation schedule.

Immunisation is carried out using the diphtheria and tetanus combined vaccine (**DT**).

The child is given one injection in the upper arm area. Post-vaccination side effects may include redness, soreness and swelling at the injection site. Less common side effects may include fever, headache, muscle pain, malaise and very rarely hives. The aforementioned symptoms usually resolve within two to three days.

General information about diphtheria and tetanus is attached.

If you have any questions, please call _____, Mon-Fri at _____ or send an email to _____

School

nurse

Parent _____
(first name and surname)

I consent: _____
(date, signature)

Diphtheria

What is diphtheria?

Diphtheria is an acute bacterial infection.

How does a person get infected with diphtheria?

The source of infection is an infected person or a carrier. The disease spreads from person to person via direct contact and droplets when coughing or sneezing. Diphtheria can be contracted at any age.

What are the symptoms of diphtheria?

The primary clinical symptoms are fever, swollen lymph nodes, sore throat, neuropathy, loss of voice, cough, respiratory failure and difficulty swallowing. Complications affect the kidneys, heart and nervous system. Paralysis of the soft palate and pneumonia may also occur. In countries with no routine vaccination, mortality from diphtheria ranges from 1.5–25%.

How to avoid infection? The most effective way to protect children against infection is timely vaccination.

In Estonia, children, adolescents and adults are routinely vaccinated against diphtheria in accordance with the immunisation schedule as follows:

Age Vaccine dose

3 months first dose

4.5 months second dose

6 months third dose

1.5–2 years fourth dose (first revaccination)

6–7 years fifth dose (second revaccination)

15–17 years sixth dose (third revaccination)

25, 35 etc every 10 years

Tetanus

What is tetanus?

Tetanus, or lockjaw, is a bacterial infection.

How does a person get infected with tetanus? Infection may occur when soil particles containing the pathogen get into a scratch or wound.

What are the symptoms of tetanus?

Tetanus is characterised by increased muscle tone and spasms. Tetanus usually starts with spasms in the jaw muscles, followed by difficulty swallowing and stiffness of the neck, shoulder and back muscles. Later, all voluntary muscles are affected. In milder cases the disease is treatable, but in more severe cases it is fatal.

How to avoid infection?

The surest method of prevention is vaccination.

In Estonia, children, adolescents and adults are routinely vaccinated against tetanus in accordance with the immunisation schedule as follows:

Age Vaccine dose

3 months first dose

4.5 months second dose

6 months third dose

1.5–2 years fourth dose (first revaccination)

6–7 years fifth dose (second revaccination)

15–17 years sixth dose (third revaccination)

25, 35 etc every 10 years

The vaccine is highly immunogenic, with post-vaccination immunity lasting at least five years.